

Catalogue of American Amphibians and Reptiles.

Powell, R. 1992. *Anolis porcatus*.

***Anolis porcatus* Gray**
Lagartija Verde, Cuban Green Anole

Anolis carolinensis: Duméril and Bibron, 1837:120 (part).

Anolis porcatus Gray, 1840:112. Type-locality, "Cuba and "Texas".
 Syntypes, British Museum of Natural History (BMNH) 1946.8.12.71, 1946.8.11.66, 1946.8.12.67-70, five adult males and a juvenile (sex unknown), date of collection unknown, presented by W.S. MacLeay (not examined by author). The first listed syntype is from "Texas," but see Distribution. Also see Remarks.
Anolis caroliniensis: Cocteau and Bibron, 1843:125 (part).

Anolis porcatus: Gray, 1845:201. *Lapsus*.

Anolis principalis var. *porcatus*: Cope, "1887" (1888):437 (part).

Anolis porcatus porcatus: Barbour, 1937:119.

Anolis carolinensis porcatus: Oliver, 1948:7.


Anolis porgatus: Otero, 1950:186. *Lapsus*.

• **Content.** No subspecies are recognized (but see Comment).

• **Definition.** *Anolis porcatus* is a medium-sized anole (males to 73 mm SVL, females smaller) characterized by a long, pointed snout, the frontal ridge higher than the canthal ridge in most males; nostril separated from the rostral by three scales; 3-4 rows of loreals; 1 scale between supraorbitals; 1-2 scales between the interparietal and supraorbital semicircles; 5 postrostrals; 4 postmentals; suboculars in contact with supralabials. Ventrals at midbody are in



Figure 1. Adult male *Anolis porcatus* from Santo Domingo, República Dominicana. Photograph by author.



Map. Distribution of *Anolis porcatus*. Solid circles represent locality records. The type-locality is too imprecise to plot.




Figure 2. Head of male *Anolis porcatius* from Camaguey, Cuba (from Ruibal and Williams, 1961. Used with permission, Museum of Comparative Zoology, Harvard University).




Figure 3. Variation in *Anolis porcatius*: A. Male from Pinar del Río, Cuba, an example of a "western" *porcatius*; B. Male from Guantánamo, Oriente, Cuba, an example of an "eastern" *porcatius*; C. Male from Camaguey, Cuba, an example of a "central" *porcatius* (from Ruibal and Williams, 1961. Used with permission, Museum of Comparative Zoology, Harvard University).

transverse and diagonal rows. Ventrals and dorsals are faintly keeled and slightly imbricate. The ear opening is circular or only slightly oval (in some specimens from Pinar del Río the posterior margin of the ear opening is V-shaped). The tail is long and round, slightly depressed at the base, and weakly verticillate, with 4 scales/verticil. Supradigital scales are multicarinate.

The ground color can change from dark brown to bright green. The male color pattern differs in eastern, central, and western populations. Eastern *porcatius* have a distinct mid-dorsal light stripe bordered by dark pigment, longitudinal dark stripes on the chin and throat, a white-bordered ocellus above the forelimbs, and evidence of reticular markings and scattered white scales on some specimens. Central populations tend to be grey to light green or brown with a complicated pattern of dark and light reticulations and numerous white scales on the nape and body. Western males show a pattern of dark reticulations over at least part of the body with an elongate dark patch above the forelimbs, single scattered white scales on the nape and parts of the body, and a white stripe below the eye. Female patterns are less geographically variable, although central females usually have distinct white scales on their sides and eastern forms (and some western specimens) have a mid-dorsal stripe similar to eastern males.

The dewlap is reddish, pink to purplish-pink, or mauve. Females lack a dewlap, but often possess indications of pink color on the throat.

• **Diagnosis.** *Anolis porcatius* may be distinguished from sympatric congeners by the long pointed snout, prominent frontal ridges higher than canthal ridges in most specimens, the rostral bordered posteriorly by five scales, circular (or only slightly oval) ear openings (with a V-shaped posterior margin in some western specimens), and a reddish dewlap in males.

• **Descriptions.** In addition to those found in references cited in the synonymy and in many of the checklists and guides under Pertinent Literature, excellent descriptions are in Ruibal and Williams (1961), Ruibal (1967), and Schwartz and Henderson (1991).

• **Illustrations.** Hand-colored engravings illustrating an adult and details of the head, feet, vent, and scales are in Cocteau and Bibron (1843). Colored illustrations are in Schwartz and Henderson (1985). Otero (1950) and Obst et al. (1988) included black and white photographs, and Molle (1961) and Vogel (1965) color photographs of adult males. Peters and Schubert (1968) provided a black and white photograph of an adult female, Parmerlee et al. (1992) of attempted male-male displacement during coitus. Molle (1956) provided a color photograph of a hatchling. Line drawings of the head are in Stejneger (1917), Barbour and Ramsden (1919), Ruibal and Williams (1961), Ruibal (1964), and Sánchez Oria and Berovides Álvarez (1987), the latter illustrating variation in supraorbital configurations. A line drawing of a lateral view of the tail is in Stejneger (1917), of the lower jaw in Oliver (1948), of the toe lamellae in Collette (1961), and of the mental scale in Pérez-Beato and Berovides (1984). Ruibal and Williams (1961) also provided line drawings of specimens representing "western," "central," and "eastern" *porcatius*. Ruibal (1967) provided a diagram of the throat display of the species. Porter et al. (1989) included photographs of the karyotype and a meiotic cell squash of males; Porter et al. (1991) illustrated a hybridized karyotype.

• **Distribution.** *Anolis porcatius* is found islandwide on Cuba and the Isla de la Juventud/Archipiélago de los Canarreos (Cayo Cantiles): Cayos de San Felipe (Cayo Real, Cayo Juan García); Archipiélago de Sabana-Camagüey (Cayo Francés, Cayo Guajaba, Cayo Santa María); Archipiélago de los Colorados (Cayo Inés de Soto); Cayo la Reina off the northern coast of Pinar del Río; and probably on many other off-shore islets and islands (Schwartz and Henderson, 1988, 1991).

This species has been introduced into Santo Domingo, República Dominicana (Hispaniola) (Arias Cornielle, 1975; Schwartz and Thomas, 1975; Haneline, 1977; Williams, 1977; Schwartz et al., 1978; Garrido and Jaime, 1984; Henderson and Schwartz, 1984; Henderson et al., 1984; Schwartz and Henderson, 1985, 1988, 1991; Powell, 1990b; Powell et al., 1990). Powell and Parmerlee (1991) reported a second, apparently disjunct Dominican colony at the Aeropuerto de las Américas; more recent field work has indicated that this population has spread to the base of the peninsula on which the airport is situated.

Reports of its presence in Texas (Gray, 1840, 1845; Boulenger, 1885) and on Key West (Allen and Slatten, 1945; Smith and Kohler, 1977) are probably erroneous (Vance, 1987). Also, green anoles, variously identified as *A. carolinensis*, *A. carolinensis porcatius*, or *A. porcatius*, have been introduced on a number of Pacific islands (see Remarks).

The Cuban distribution has been illustrated in Ruibal and Williams (1961), the Hispaniolan range in Powell et al. (1990), the total West Indian distribution in Schwartz and Henderson (1991).

• **Fossil Record.** None.

• **Pertinent Literature.** The species is included in the checklists and guides of Gundlach (1880), Cope ("1887" [1888]), Stejneger (1917), Barbour and Ramsden (1916, 1919), Barbour (1916, 1930a, b, 1935, 1937), Alayo Dalmau (1951, 1955), Ruibal (1964), Buide (1967), Garrido and Schwartz (1968, 1969), Lando and Williams (1969), Schwartz and Thomas (1975), Schwartz et al. (1978), Garrido and Jaume (1984), Henderson and Schwartz (1984), Henderson et al. (1984), Schwartz and Henderson (1985, 1988, 1991), and Banks et al. (1987). Additional distributional records are in Sutcliffe (1952), Schwartz and Ogren (1956), Garrido (1973), Estrada et al. (1987), and Abreu et al. (1989). SEA/DVS (1990) provided an index of habitats in the Dominican Republic.

Other topics addressed in the literature include: ecology (Ruibal, 1958, 1961; Collette, 1961; Lando and Williams, 1969; Peters, 1970; Schoener, 1970; Arias Cornielle, 1975; Estrada and Novo Rodríguez, 1986), ecomorphology (Estrada and Silva Rodríguez, 1984; Schoener, 1988), species recognition (Williams and Rand, 1977), food habits (Otero, 1950; Alayón García, 1976), color changes (Hadley, 1929), behavior (Ruibal, 1967; Powell, 1990a; Parmelee et al., 1992), reproduction (Dunn, 1926; Rand, 1967), development (Pérez-Beato, 1982), parasitism (Barus and Coy Otero, 1969a, b; Coy Otero, 1970; Coy Otero and Barus, 1973, 1979; Coy Otero and Lorenzo Hernández, 1982; Baker, 1987), geographic variation (Pérez-Beato and Berovides, 1979, 1982, 1984; Espinosa Lopez et al., 1985; Sánchez Oria and Berovides Álvarez, 1987), karyotype (Gorman and Atkins, 1968; Porter et al., 1989), the evolutionary implication of the location of ribosomal DNA (Porter et al., 1991), relationship between variability of hepatic esterases and the degree of environmental stability (Mugica Valdés et al., 1982), general relationships (Barbour, 1928; Burnell and Hedges, 1990), geographic variation, relationship, and hybridization with *A. allisoni* (Ruibal and Williams, 1961), niche displacement by *A. allisoni* (Peters and Schubert, 1968); placement in the alpha group of anoles based on osteology (Etheridge, 1960), formal placement in the *carolinensis* superspecies (Williams, 1976), electrophoretic distinction from *A. carolinensis* (Buth et al., 1980), morphological distinction from and relationship with *A. carolinensis* (Garman, "1887" [1888]; Oliver, 1948; Williams, 1969), distinction from *A. brunneus* (Barbour, 1914), human exploitation (Barbour, 1937), care and reproduction in captivity (Molle, 1956, 1961; Rover, 1960; Markert, 1980), English common name (Collins et al., 1982; Collins, 1990).

• **Remarks.** Boulenger (1885) listed one male from Texas and five "males & yg." from Cuba, all from W.S. MacLeay, Esq., as the types for *Anolis porcatius*. However, data currently on file (Colin McCarthy, in litt. 09.VII.90) differ. BMNH 1946.8.11.66 is an adult male from "Texas," collector unknown; BMNH 1946.8.11.66 is an adult male presented by W.S. MacLeay; and BMNH 1946.8.12.67-.70 are three adult males and a juvenile from Mr. Leadbeater's collection. These discrepancies (see also Russell and Bauer, 1991) illustrate the kinds of problems in quality control of data inherent to natural history collections over time, and argue for the provision of adequate resources to maintain the accuracy of irreplaceable information.

Green anoles, variously identified as *A. carolinensis*, *A. carolinensis porcatius*, or *A. porcatius*, have been introduced to various Pacific islands, including the Hawaiian Islands, Guam, Saipan, and Chichi-Jima, Japan (Shaw and Breese, 1951; Oliver and Shaw, 1953; Ruibal, 1964; Hunsaker and Breese, 1967; Cochran and Goin, 1970; Anonymous, 1972; Flint, 1972; Smith and Kohler, 1977; Williams, 1977; McKeown, 1978; Jones, 1979; Matsumoto et al., 1980; Wilson and Porras, 1983; Hara, 1986; Hasegawa, 1986; Chan et al., 1987; Vance, 1987; Hasegawa et al., 1988; Schwartz and Henderson, 1988; Collins, 1990; Powell et al., 1990). Contrary to Vance's (1987) statements, however, these populations apparently all represent *A. carolinensis* (Hasegawa et al., 1988; Lazell, 1989; R.I. Crombie, in litt. 14 VI 90; G.C. Mayer, in litt. 3 VII 90; E.E. Williams, pers. comm.).

• **Etymology.** The specific epithet *porcatius* is presumably derived from the Latin *porca* (= ridge between two furrows) in reference to the pronounced frontal ridges on the heads of males of this species.

• **Comment.** Ruibal (1964) suggested that *A. porcatius* may represent more than one species.

Literature Cited

- Abreu, R.M., J. de la Cruz, A. Rams, and M.E. Garcia. 1989. Vertebrado del complejo montañoso "La Zoilita", Holguín, Cuba. Poeyana (370):1-16.
- Alayo Dalmau, P. 1951. Especies herpetológicas halladas en Santiago de Cuba. Bol. Hist. Nat. Soc. Habana 2:106-111.
- . 1955. Lista de los reptiles de Cuba. Mimeographed. Univ. de Oriente, Mus. Charles T. Ramsden, Santiago de Cuba.
- Alayón García, G. 1976. Araneidos depredados por anolinos. Misc. Zool. (2):1.
- Allen, E.R. and R. Slatten. 1945. A herpetological collection from the vicinity of Key West, Florida. Herpetologica 3:25-26.
- Anon. 1972. Herpetological species introduced onto Hawaii. Herpetol. Rev. 4:172.
- Arias Cornielle, Y.A. 1975. *Anolis chlorocyanus* y *Anolis porcatius* (Sauria, Iguanidae) en la ciudad de Santo Domingo. Tesis. Univ. Auton. Santo Domingo.
- Baker, M.R. 1987. Synopsis of the Nematoda parasitic in amphibians and reptiles. Occ. Paps. Biol. Mem. Univ. Newfoundland (11):1-325.
- Banks, R.C., R.W. McDiarmid, and A.L. Gardner (eds.). 1987. Checklist of vertebrates of the United States, the U.S. territories, and Canada. U.S. Dept. Interior, Fish Wildl. Serv. Resour. Publ. (166): ii + 79 p.
- Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with especial reference to amphibians and reptiles. Mem. Mus. Comp. Zool. 44:205-360.
- . 1916. The reptiles and amphibians of the Isle of Pines. Ann. Carnegie Mus. 10:297-308.
- . 1928. Reptiles from the Bay Islands. Proc. New England Zool. Club 10:55-61.
- . 1930a. The anoles. I. The forms known to occur on the neotropical islands. Bull. Mus. Comp. Zool. 70:105-144.
- . 1930b. A list of Antillean reptiles and amphibians. Zoologica 11:61-116.
- . 1935. A second list of Antillean reptiles and amphibians. Zoologica 19:77-141.
- . 1937. Third list of Antillean reptiles and amphibians. Bull. Mus. Comp. Zool. 82:75-166.
- and C.T. Ramsden. 1916. Catálogo de los reptiles y anfibios de la isla de Cuba. Mem. Soc. Poeyana 2:124-143.
- and —. 1919. The herpetology of Cuba. Mem. Mus. Comp. Zool. 47:69-213.
- Barus, V. and A. Coy Otero. 1969a. Nematodos del género *Parapharyngodon* Chatterji, 1933 (Oxyuridae), en Cuba. Torreia (n.s.) (7):1-10.
- and —. 1969b. Systematic survey of nematodes parasitizing lizards (Sauria) in Cuba. Helminthologia 10:329-346.
- Boulenger, G.A. 1885. Catalogue of the lizards in the British Museum (Natural History). 2nd ed. Vol. 2. Taylor and Francis, London.
- Buide, M.S. 1967. Lista de los anfibios y reptiles de Cuba. Torreia (n.s.) (1):5-60.
- Burnell, K.L. and S.B. Hedges. 1990. Relationships of West Indian *Anolis* (Sauria: Iguanidae): an approach using slow-evolving protein loci. Carib. J. Sci. 26:7-30.
- Buth, D.G., G.C. Gorman, and C.S. Lieb. 1980. Genetic divergence between *Anolis carolinensis* and its Cuban progenitor, *Anolis porcatius*. J. Herpetol. 14:279-284.
- Chan, J.G., T. Moniz, and M. Kealamakia. 1987. Geographic distribution: *Anolis carolinensis porcatius*. Herpetol. Rev. 18:40.
- Cochran, D.M. and C.J. Goin. 1970. The new field book of reptiles and amphibians. G.P. Putnam's Sons, New York.
- Cocteau, J.T. and G. Bibron. 1843. Reptiles. In R. de la Sagra, Histoire physique, politique et naturelle de l'île de Cuba, pt. 7. Arthus Bertrand, Paris.
- Collette, B.B. 1961. Correlations between ecology and morphology in anoline lizards from Havana, Cuba and southern Florida. Bull. Mus. Comp. Zool. 125:137-162.
- Collins, J.T. 1990. Standard common and current scientific names for

- North American amphibians and reptiles. 3rd ed. Soc. Study Amphib. Rept. Herpetol. Circ. (19):iii + 41 p.
- , R. Conant, J.E. Huheey, J.L. Knight, E.M. Rundquist, and H.M. Smith. 1982. Standard common and current scientific names for North American amphibians and reptiles. 2nd ed. Soc. Study Amphib. Rept. Herpetol. Circ. (12):iv + 28 p.
- Cope, E.D. "1887" (1888). List of the Batrachia and Reptilia on the Bahama Islands. Proc. U.S. Natl. Mus. 10:436-439.
- Coy Otero, A. 1970. Contribución al conocimiento de la helmintofauna de los saurios Cubanos. Ciencias Ser. 4 Ciencias Biol. (4): 1-50.
- and V. Barus. 1973. Notes on nematodes of the genus *Cyrtosomum* (Atractidae) parasitic in Cuban lizards (Sauria). Folia Parasit. 20:297-305.
- and —. 1979. Nematodes parasitizing Cuban reptiles. Acta Sci. Nat. Brno 13:1-43.
- and N. Lorenzo Hernández. 1982. Lista de los helmintos parásitos de los vertebrados silvestres cubanos. Poeyana (235):1-57.
- Duméril, A.M.C. and G. Bibron. 1837. Erpétologie générale ou histoire naturelle complète des reptiles. Paris, 4:1-571.
- Dunn, E.R. 1926. Notes on Cuban anoles. Copeia (157):153-154.
- Espinosa Lopez, G., C. Gonzalez Suarez, and V. Berovides Álvarez. 1985. Albumina de suero y esterases de musculo en poblaciones de *Anolis porcatus* Gray 1840 del occidente de Cuba. Carib. J. Sci. 21:55-62.
- Estrada, A.R., G. Alayón, A. Pérez Asso, C. Peña, and E. Solana. 1987. Lista preliminar de anfibios y reptiles de las Cuchillas de Moa y Toa, Cuba. Garciana 8:3-4.
- and J. Novo Rodríguez. 1986. Subnicho estructural de *Anolis sagrei* en Cayo Inés de Soto, Cuba. Análisis intra- y extrapoblacional. Poeyana (320):1-13.
- and A. Silva Rodríguez. 1984. Análisis de la ecomorfología de 23 especies de lagartos cubanos del género *Anolis*. Cienc. Biol. 12: 91-104.
- Etheridge, R. 1960. The relationships of the anoles (Reptilia: Sauria: Iguanidae): an interpretation based on skeletal morphology. Ph.D. Diss., Univ. Michigan, Ann Arbor.
- Flint, C. 1972. The reptiles and amphibians of the Hawaiian Islands. Herpetology 6:8-11.
- Garman, S. "1887" (1888). On West Indian reptiles. Iguanidae. Bull. Essex Inst. 19:25-50.
- Garrido, O.H. 1973. Anfibios, reptiles y aves de Cayo Real (Cayos de San Felipe), Cuba. Poeyana (119):1-50.
- and M. Jaume. 1984. Catálogo descriptivo de los anfibios y reptiles de Cuba. Doñana, Acta Vertebrata 11:5-128.
- and A. Schwartz. 1968. Anfibios, reptiles y aves de la península de Guanahacabibes, Cuba. Poeyana, Ser. A (53):1-68.
- and —. 1969. Anfibios, reptiles y aves de Cayo Cantiles. Poeyana (67):1-44.
- Gorman, G.C. and L. Atkins. 1968. New karyotypic data for 16 species of *Anolis* (Sauria: Iguanidae) from Cuba, Jamaica, and the Cayman Islands. Herpetologica 24:13-21.
- Gray, J.E. 1840. Catalogue of the species of reptiles collected in Cuba by W.S. MacLeay, Esq.; - with some notes of their habits extracted from his MS. Ann. Mag. Nat. Hist., Ser. 1, 5:108-115.
- . 1845. Catalogue of the specimens of lizards in the collection of the British Museum. E. Newman, London.
- Gundlach, J. 1880. Contribución a la erpetología cubana. G. Monteil, Habana.
- Hadley, C.E. 1929. Color changes in two Cuban lizards. Bull. Mus. Comp. Zool. 69:105-114.
- Haneline, P.G. 1977. Geographic distribution: *Anolis porcatus*. Herpetol. Rev. 8:39.
- Hara, K. 1986. Kika Hachuuri (The naturalized reptiles in Japan). Iden (The Heredity) 40:14-17 (in Japanese).
- Hasegawa, M. 1986. Ecological studies of the lizards on Chichi-jima, the Bonin Islands. Ogasawara Kenkyu Nenpo (9):12-17 (in Japanese).
- , T. Kusano, and K. Miyashita. 1988. Range expansion of *Anolis c. carolinensis* on Chichi-jima, the Bonin Islands, Japan. Japanese J. Herpetol. 12:115-118.
- Henderson, R.W. and A. Schwartz. 1984. A guide to the identification of the amphibians and reptiles of Hispaniola. Milwaukee Pub. Mus. Spec. Publ. Biol. Geol. (4):1-70.
- , —, and S.J. Incháustegui. 1984. Guía para la indentificación de los anfibios y reptiles de la Hispaniola. Mus. Nac. Hist. Nat. Ser. Mono. (1):1-128.
- Hunsaker, D. II and P. Breese. 1967. Herpetofauna of the Hawaiian Islands. Pacific Sci. 21:423-428.
- Jones, R. 1979. Hawaiian lizards - their past, present, and future. Bull. Maryland Herpetol. Soc. 15:37-45.
- Lando, R.V. and E.E. Williams. 1969. Notes on the herpetology of the U.S. naval base at Guantanamo Bay, Cuba. Stud. Fauna Curaçao Carib. Isl. 31:159-201.
- Lazell, J.D. Jr. 1989. Kaho'olawe: Hawaii's largest uninhabited island. Explorers J. 67:123-126.
- Markert, T. 1980. Pflege und Nachzucht bei *Anolis porcatus*, dem Leistenanolis. Aquar.-Terrar. Z. 33:426-427.
- Matsumoto, Y., T. Matsumoto, and K. Miyashita. 1980. Herpetological survey on Chichi-jima and Haha-jima, the Bonin Islands. Ogasawarashoto Sizenkankyo Genkyochousa Houkokusho Tokyo (1):65-75 (in Japanese).
- McKeown, S. 1978. Hawaiian reptiles and amphibians. Oriental Publ. Co., Honolulu.
- Molle, F. 1956. Erfolgreiche Aufzucht von *Anolis porcatus* Gray und *Anolis carolinensis* Voigt in Gefangenschaft. Aquar. Terrar. Z. 9:300-304.
- . 1961. 7 Jahre Anoliszucht - Erfahrungen und Erkenntnisse II. Aquar. Terrar. Z. 14:181-183.
- Mugica Valdés, L., G. Espinosa López, and V. Berovides Álvarez. 1982. Patrones electroforéticos de esterases hepáticas en siete especies lagartos del género *Anolis*. Cienc. Biol. (8):37-48.
- Obst, F.J., K. Richter, and U. Jacob. 1988. The completely illustrated atlas of reptiles and amphibians for the terrarium. T.F.H. Publ., Inc., Neptune City, New Jersey.
- Oliver, J.A. 1948. The anoline lizards of Bimini, Bahamas. Amer. Mus. Nov. (1383):1-36.
- and C.E. Shaw. 1953. The amphibians and reptiles of the Hawaiian Islands. Zoologica 38:65-95.
- Otero, A.R. 1950. Sobre la alimentacion de *Anolis porcatus*. Bol. His. Nat. 1(4):186-187.
- Parmerlee, J.S. Jr., R. Powell, D.D. Smith, and A. Lathrop. 1992. Unusual behavior in the Cuban Green Anole, *Anolis porcatus* (Sauria: Polychridae). Bull. Chicago Herpetol. Soc. 27:118.
- Pérez-Beato, O. 1982. Algunos indicadores en el desarrollo de *Anolis porcatus* (Sauria: Iguanidae). Cienc. Biol. (7):127-129.
- and V. Berovides. 1979. Diferenciación de *Anolis porcatus* (Sauria: Iguanidae) en el Occidente de Cuba. Res. Seg. Even. Cien. Univ. La Habana.
- and —. 1982. Estudio electroforético de las proteínas del plasma en dos especies de *Anolis* (Sauria: Iguanidae). Poeyana (242):1-7.
- and —. 1984. Índice cefálico y patrón de placa mental: dos elementos fenotípicos en la diferenciación de *Anolis porcatus* (Sauria: Iguanidae) en el occidente de Cuba. Poeyana (262):1-8.
- Peters, G. 1970. Zur Taxonomie und Zoogeographie der kubanischen anolinen Eidechsen (Reptilia, Iguanidae). Mitt. Zool. Mus. Berlin 46:197-234.
- and R. Schubert. 1968. Theoretische und praktische Aspekte einer biologischen Expedition durch Kuba. Biologie in der Schule (8/9):337-346.
- Porter, C.A., R.I. Crombie, and R.J. Baker. 1989. Karyotypes of five species of Cuban lizards. Occ. Pap. Mus. Texas Tech Univ. (130): 1-6.
- , M.J. Hamilton, J.W. Sites, Jr., and R.J. Baker. 1991. Location of ribosomal DNA in chromosomes of squamate reptiles: systematic and evolutionary implications. Herpetologica 47:271-280.
- Powell, R. 1990a. Life history notes: *Anolis porcatus*. Behavior. Herpetol. Rev. 21:93.
- . 1990b. Geographic distribution: *Anolis porcatus*. Herpetol. Rev. 21:96.
- and J.S. Parmerlee, Jr. 1991. Geographic distribution: *Anolis porcatus*. Herpetol. Rev. 22:65.
- , D.D. Smith, J.S. Parmerlee, Jr., C.V. Taylor, and M.L. Jolley. 1990. Range expansion by an introduced anole: *Anolis porcatus* in the Dominican Republic. Amphib.-Rept. 11:421-425.
- Rand, A.S. 1967. Communal egg laying in anoline lizards. Herpetologica 23:227-230.
- Rover, W. De. 1960. Het kweken van reptielen. Aquarium (Den Haag) 30:222-223.
- Ruibal, R. 1958. A preliminary investigation of the ecology and taxonomy of Cuban lizards. Amer. Phil. Soc. Yr. Bk. 1957:256-258.
- . 1961. Thermal relations of five species of tropical lizards. Evolution 15:98-111.
- . 1964. An annotated checklist and key to the anoline lizards of

- Cuba. Bull. Mus. Comp. Zool. 130:473-520.
- . 1967. Evolution and behavior in West Indian anoles, p. 116-140. In W.W. Millstead (ed.). Lizard ecology: a symposium. Univ. Missouri Press, Columbia.
- and E.E. Williams. 1961. Two sympatric Cuban anoles of the *carolinensis* group. Bull. Mus. Comp. Zool. 125:183-208.
- Russell, A.P. and A.M. Bauer. 1991. *Anolis grabami*. Cat. Amer. Amphib. Rept.:514.1-514.4.
- Sánchez Oria, B. and V. Berovides Álvarez. 1987. Variación y selección natural en *Anolis porcatius* Gray, 1840 (Sauria: Iguanidae). Cienc. Biol. 18:111-118.
- Schoener, T.W. 1970. Size patterns in West Indian *Anolis* lizards. II. Correlations with the sizes of particular sympatric species - displacement and convergence. Amer. Nat. 104:155-174.
- . 1988. Testing for non-randomness in sizes and habitats of West Indian lizards: choice of species pool affects conclusions from null models. Evol. Ecol. 2:1-26.
- Schwartz, A. and R.W. Henderson. 1985. A guide to the identification of the amphibians and reptiles of the West Indies exclusive of Hispaniola. Milwaukee Pub. Mus., Milwaukee, Wisconsin.
- and ———. 1988. West Indian amphibians and reptiles: a check-list. Milwaukee Pub. Mus. Contrib. Biol. Geol. (74):1-264.
- and ———. 1991. Amphibians and reptiles of the West Indies: descriptions, distributions, and natural history. Univ. Florida Press, Gainesville.
- and L.H. Ogren. 1956. A collection of reptiles and amphibians from Cuba, with the descriptions of two new forms. Herpetologica 12:91-110.
- and R. Thomas. 1975. A check-list of West Indian amphibians and reptiles. Carnegie Mus. Nat. Hist., Spec. Publ. (1):1-216.
- , ———, and L.D. Ober. 1978. First supplement to a check-list of West Indian amphibians and reptiles. Carnegie Mus. Nat. Hist., Spec. Publ. (5):1-35.
- SEA/DVS. 1990. La diversidad biológica en la República Dominicana: reporte preparado por el Departamento de Vida Silvestre para el Servicio Alemán de Cooperación Social-Técnica y Fondo Mundial para la Naturaleza (WWF-US). Apéndice. Sec. Estado Agric., SURENA/DVS. Sto. Domingo, República Dominicana.
- Shaw, C.E. and P.L. Breese. 1951. An addition to the herpetofauna of Hawaii. Herpetologica 7:68.
- Smith, H.M. and A.J. Kohler. 1977. A survey of herpetological introductions in the United States and Canada. Trans. Kansas Acad. Sci. 80:1-24.
- Stejneger, L. 1917. Cuban amphibians and reptiles collected for the United States National Museum from 1899 to 1902. Proc. U. S. Natl. Mus. 53:259-291.
- Sutcliffe, R. 1952. Results of the Catherwood-Chaplin West Indies expedition, 1948. Part VI. Amphibia and Reptilia. Notul. Nat. (243): 1-8.
- Vance, T. 1987. The Cuban green anole (*Anolis porcatius*): a colonizing species. Bull. Maryland Herpetol. Soc. 23:105-108.
- Vogel, Z. 1965. Herpetologische Beobachtungen auf Kuba. Aquar.-Terrar. 12:235-239.
- Williams, E.E. 1969. The ecology of colonization as seen in the zoogeography of anoline lizards on small islands. Quart. Rev. Biol. 44:345-389.
- . 1976. West Indian anoles: a taxonomic and evolutionary summary. I. Introduction and a species list. Breviora (440):1-21.
- . 1977. Anoles out of place: introduced anoles. The Third *Anolis* Newsl.:110-118.
- and A.S. Rand. 1977. Species recognition, dewlap function and faunal size. Amer. Zool. 17:261-270.
- Wilson, L.D. and L. Porras. 1983. The ecological impact of man on the south Florida herpetofauna. Univ. Kansas Mus. Nat. Hist., Spec. Publ. (9):1-89.

Robert Powell, Department of Natural Sciences, Avila College, Kansas City, MO 64145.

Primary editor for this account, Andrew H. Price.

Published 30 November 1992 and Copyright © 1992 by the Society for the Study of Amphibians and Reptiles.
